

ON THE SYMMETRY OF ARITHMETICAL FUNCTIONS IN ALMOST ALL SHORT INTERVALS, IV

by
G.Coppola

Abstract. We study the arithmetic (real) function $f = g * \mathbf{1}$, with g “essentially bounded” and supported over the integers of $[1, Q]$. In particular, we obtain non-trivial bounds, through f “correlations”, for the “Selberg integral” and the “symmetry integral” of f in almost all short intervals $[x - h, x + h]$, $N \leq x \leq 2N$, beyond the “classical” level, up to a very high level of distribution (for h not too small). This time we go beyond Large Sieve inequality [C]. Precisely, our method applies Weil bound for Kloosterman sums.

1. Introduction and statement of the results.

This paper pursues the study of the symmetry integral (and, also, of the Selberg integral) of arithmetical functions, started in our paper [C], making a substantial progress on the level of distribution of the (sieve) arithmetical functions involved.

We will study ARITHMETICAL FUNCTIONS $f : \mathbb{N} \rightarrow \mathbb{R}$ in almost all short intervals; here, “ALMOST ALL” $[x - h, x + h]$ with $N < x \leq 2N$ means all, except for at most $o(N)$ of them; where $[x - h, x + h]$ “SHORT INTERVAL” means $h = h(N)$ (for $N < x \leq 2N$, see above), with (as $N \rightarrow \infty$) $h \rightarrow \infty$ and $h = o(N)$. Here we’ll follow Landau’s notation: $F(N) = o(G(N))$ stands for (when $N \rightarrow \infty$) $F(N)/G(N) \rightarrow 0$ and $F(N) = \mathcal{O}(G(N))$ (or the equivalent Vinogradov notation $F(N) \ll G(N)$) abbreviates (again, $N \rightarrow \infty$) $|F(N)|/G(N)$ is bounded (above, by a positive constant, called the $\mathcal{O}$ -constant or the $\ll$ -constant); $F(N) \ll_{A,B,C} G(N)$ (or $F(N) = \mathcal{O}_{A,B,C}(G(N))$) indicates the dependence of the $\ll$ - (or $\mathcal{O}$ -) constant on these parameters. As usual, we will indicate the greatest common divisor (GCD) of a and b as: (a, b) .

We will use two important quantities (describing f or h , respectively), namely the (DISTRIBUTION) LEVEL λ of f and the WIDTH θ of the short interval. Writing $f(n) = \sum_{d|n} g(d)$, we will let $\lambda \stackrel{\text{def}}{=} (\log Q)/(\log N)$, whenever $g(q) = 0 \forall q > Q$. And, given $h = h(N)$ the length of our short interval(s), set $\theta \stackrel{\text{def}}{=} (\log h)/(\log N)$. In this paper the width will be assumed to be $0 < \theta < 1$ (but “far” from edges).

As usual, we’ll write $\mathbf{1}(n) = 1$, i.e. the characteristic function of natural numbers (if no subscripts; while we write $\mathbf{1}_\varphi$ to mean 1 if φ is true, 0 otherwise), and the above becomes $f = g * \mathbf{1}$, where $*$ is Dirichlet product [T]; from Möbius inversion, see [T], $g = f * \mu$ (where μ is Möbius function, $\mu(1) \stackrel{\text{def}}{=} 1$, $\mu(n) \stackrel{\text{def}}{=} (-1)^r$ when n is the product of r distinct primes, $\mu(n) \stackrel{\text{def}}{=} 0$ in all other cases, see [T]).

Some authors, like Aurel Wintner, called $f * \mu$, say f' , the “Eratosthenes transform” of f .

We call f ESSENTIALLY BOUNDED if $\forall \varepsilon > 0$ we have $f(n) \ll n^\varepsilon$ (as $n \rightarrow \infty$); for example, the number of (positive) divisors of n , say $d(n) \stackrel{\text{def}}{=} \sum_{d|n} 1$, is such (like many other interesting arithmetical functions). Sometimes (especially during the proofs of the Lemma and of our Theorem) we’ll abbreviate $A \ll B$ to mean: $\forall \varepsilon > 0 \ A \ll_\varepsilon N^\varepsilon B$. Also, Möbius inversion easily allows to prove that $f \ll 1$ if, and only if, $g \ll 1$ (whenever $f = g * \mathbf{1}$). In the sequel g support will be $[Q, 2Q]$ or $[1, Q]$ (no difference, using a dissection argument: additional logarithms are absorbed into N^ε): our arguments won’t be affected (for small Q , [C]).

The quantity checking f “almost all”-symmetry in short intervals is the SYMMETRY INTEGRAL

$$I_f(N, h) \stackrel{\text{def}}{=} \int_N^{2N} \left| \sum_{|n-x| \leq h} f(n) \operatorname{sgn}(n-x) \right|^2 dx,$$

where $\operatorname{sgn}(0) \stackrel{\text{def}}{=} 0$, $\operatorname{sgn}(r) \stackrel{\text{def}}{=} r/|r|$ ($\forall r \neq 0$). (For the definition of J_f , the Selberg integral of f , see [C].) In our previous paper [C] we obtained non-trivial bounds for I_f (and J_f), whenever the level is $\lambda < \frac{1+\theta}{2}$. We mean, for non-trivial, bounds of the kind $I_f(N, h) \ll \frac{Nh^2}{N^\delta}$ (same for J_f), for some $\delta > 0$ (small).

Here we will supersede these results, getting level $\lambda < \max(1 - \frac{\theta}{2}, \frac{1+\theta}{2})$ (using previous, too).

As usual, $[\alpha] \stackrel{\text{def}}{=} \min_{n \in \mathbb{Z}, n \leq \alpha} n$ will be the INTEGER PART of $\alpha \in \mathbb{R}$ and $\{\alpha\} \stackrel{\text{def}}{=} \alpha - [\alpha]$ its FRACTIONAL PART.

We give, now, our main result.

THEOREM. *Fix $\theta_0 \in]0, 1/2[$. Let $N, h \in \mathbb{N}$, where $N^\theta \ll h \ll N^\theta$, if $N \rightarrow \infty$, with $\theta_0 < \theta < 1 - \theta_0$. Assume $f = g * \mathbf{1}$ real and essentially bounded with, say, $g(q) = 0, \forall q > Q, Q = o(\frac{N}{\sqrt{h}})$. Then*

$$I_f(N, h) \ll N h + h^3 + N h^2 \frac{h}{Q} + N h^2 \left(\frac{Q^2 h}{N^2} \right)^{1/5}.$$

(We point out that a result similar to our Theorem holds for the Selberg integral of f , too.)

We explicitly remark that an important limitation to obtain non-trivial bounds comes from $h = o(Q)$, but can be avoided (namely, $Q \ll h$ gives $\ll h^3$, since $\lambda < \frac{1+\theta}{2}$ from quoted LS-type result of [C]).

As regards the level, it comes from our Theorem the quoted above $\lambda < 1 - \theta/2$.

The paper is organized as follows:

- ★ in section 2 we prove the (exponential sums) Lemma for the proof of our Theorem,
- ★ that is given in section 3.

2. A trigonometric Lemma.

Lemma. Let $a, j, t \in \mathbb{N}$ with $(a, t) = 1$ and $a = o(t)$, as $a \rightarrow \infty, t \rightarrow \infty$ ($a, t \ll N$, if $N \rightarrow \infty$). Then

$$\sum_{0 < |k| \leq \frac{t}{2}} \left[\cos \frac{2\pi k \bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k \bar{t}}{a} \sin \frac{2\pi k}{at} \right] \cos \frac{2\pi j k}{t} = \sum_{r \in \mathbb{Z}_a} \cos \frac{2\pi r \bar{t}}{a} \Sigma_1 + \sum_{r \in \mathbb{Z}_a} \sin \frac{2\pi r \bar{t}}{a} \Sigma_2,$$

defining the reciprocal $\bar{a}a \equiv 1 \pmod{t}$, we have $\sum_r \cos \frac{2\pi r \bar{t}}{a} \Sigma_1(0) \ll \frac{1}{a}$ and $\sum_r \sin \frac{2\pi r \bar{t}}{a} \Sigma_2(0) \ll 1$, where

$$(1) \quad \Sigma_1 = \Sigma_1(j) \stackrel{\text{def}}{=} 2 \sum_{\substack{k \leq t/2 \\ k \equiv r(a)}} \left(\cos \frac{2\pi k}{at} - 1 \right) \cos \frac{2\pi j k}{t} \Rightarrow \frac{1}{t} \sum_{0 \leq |j| \leq \frac{t}{2}} \left| \sum_{0 \leq |r| \leq \frac{a}{2}} \cos \frac{2\pi r \bar{t}}{a} \Sigma_1(j) \right| \ll \frac{1}{a},$$

(we abbreviate $k \equiv r(a)$ for $k \equiv r \pmod{a}$, hereon), indicating $M(j) \stackrel{\text{def}}{=} -2 \sin \frac{2\pi j r}{t} \sum_{J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \sin \frac{2\pi j a J}{t}$,

$$(2) \quad \Sigma_2 = \Sigma_2(j) \stackrel{\text{def}}{=} 2 \sum_{\substack{k \leq t/2 \\ k \equiv r(a)}} \sin \frac{2\pi k}{at} \cos \frac{2\pi j k}{t} \Rightarrow \frac{1}{t} \sum_{0 \leq |j| \leq \frac{t}{2}} \left| \sum_{0 \leq |r| \leq \frac{a}{2}} \sin \frac{2\pi r \bar{t}}{a} (\Sigma_2(j) - M(j)) \right| \ll \frac{a}{t} + \frac{1}{a}.$$

Proof. Use $k = Ja + r$ for (both $J = 0$, see the remainder, and) $J \geq 1$:

$$\begin{aligned} \Sigma_1 &= 2 \sum_{J \leq \frac{t}{2a} - \frac{r}{a}} \left(\cos \left(\frac{2\pi J}{t} + \frac{2\pi r}{at} \right) - 1 \right) \cos \frac{2\pi j(Ja + r)}{t} + \mathcal{O} \left(\frac{1}{t^2} \right) \\ &= 2 \sum_{J \leq \frac{t}{2a}} \left(\cos \frac{2\pi J}{t} - 1 \right) \cos \frac{2\pi j(Ja + r)}{t} + \mathcal{O} \left(\frac{1}{t^2} + \frac{1}{at} + \frac{1}{a^2} \right) \end{aligned}$$

where, due to $a = o(t)$, these remainders contribute $\ll \frac{1}{a}$ to (1). Simplifying r -odd terms, we get

$$\begin{aligned} &\sum_{r \in \mathbb{Z}_a} \cos \frac{2\pi r \bar{t}}{a} \left(2 \sum_{J \leq \frac{t}{2a}} \left(\cos \frac{2\pi J}{t} - 1 \right) \cos \frac{2\pi j(Ja + r)}{t} \right) \\ &= \sum_{r \in \mathbb{Z}_a} \cos \frac{2\pi r \bar{t}}{a} \cos \frac{2\pi j r}{t} \left(2 \sum_{J \leq \frac{t}{2a}} \left(\cos \frac{2\pi J}{t} - 1 \right) \cos \frac{2\pi j a J}{t} \right) \ll \frac{1}{a} \left(\mathbf{1}_{j \neq 0} \frac{1}{\left\| \frac{ja}{t} \right\|} \right), \end{aligned}$$

hereon $\|\alpha\| \stackrel{\text{def}}{=} \min_{n \in \mathbb{Z}} |\alpha - n|$, applying partial summation [T] to $\sum_J e(J\alpha) \ll \frac{1}{\|\alpha\|}$, see [D, chap.26] ($\alpha = \frac{ja}{t}$);

$$\frac{1}{t} \sum_{j \neq 0} \frac{1}{\left\| \frac{ja}{t} \right\|} = \sum_{n \leq t/2} \frac{1}{n} \sum_{j \equiv \pm \bar{a}n(t)} 1 = 2 \sum_{n \leq \frac{t}{2}} \frac{1}{n} \ll \log(t+2) \ll 1.$$

Then, (1) is proved.

In order to prove (2), we first let $k = Ja + r$ (again, for $J = 0$, see the remainder):

$$\Sigma_2(j) = \mathcal{O} \left(\frac{1}{t} \right) + 2 \sum_{J \leq \frac{t}{2a} - \frac{r}{a}} \sin \left(\frac{2\pi J}{t} + \frac{2\pi r}{at} \right) \cos \left(\frac{2\pi j a J}{t} + \frac{2\pi j r}{t} \right)$$

We exclude $j \equiv \pm \bar{a}(t)$ in the following.

We sweep out, this time, after a small correction like in the previous Lemma, the r -even terms,

$$\sum_{0 < |r| \leq a/2} \sin \frac{2\pi r \bar{t}}{a} (\Sigma_2 - M(j)) = 2 \sum_{0 < |r| \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sum_{J \leq \frac{t}{2a}} \sin \left(\frac{2\pi J}{t} + \frac{2\pi r}{at} \right) \cos \left(\frac{2\pi j a J}{t} + \frac{2\pi j r}{t} \right)$$

$$\begin{aligned}
& +\mathcal{O}\left(\frac{a}{t}\right) - 2 \sum_{r \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sum_{\frac{t}{2a} < J \leq \frac{t}{2a} + \frac{r}{a}} \sin \left(\frac{2\pi J}{t} - \frac{2\pi r}{at} \right) \cos \left(\frac{2\pi jaJ}{t} - \frac{2\pi jr}{t} \right) \\
& - 2 \sum_{r \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sum_{\frac{t}{2a} - \frac{r}{a} < J \leq \frac{t}{2a}} \sin \left(\frac{2\pi J}{t} + \frac{2\pi r}{at} \right) \cos \left(\frac{2\pi jaJ}{t} + \frac{2\pi jr}{t} \right) \\
& + 2 \sum_{0 < |r| \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sin \frac{2\pi jr}{t} \sum_{J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \sin \frac{2\pi jaJ}{t} \\
& = \mathcal{O}\left(\frac{a}{t}\right) - 2 \left(\sum_{0 < |r| \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \left(\cos \frac{2\pi r}{at} - 1 \right) \sin \frac{2\pi jr}{t} \right) \left(\sum_{J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \sin \frac{2\pi jaJ}{t} \right) \\
& + 2 \left(\sum_{0 < |r| \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sin \frac{2\pi r}{at} \cos \frac{2\pi jr}{t} \right) \left(\sum_{J \leq \frac{t}{2a}} \cos \frac{2\pi J}{t} \cos \frac{2\pi jaJ}{t} \right) \\
& - 2 \sum_{r \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sum_{\frac{t}{2a} < J \leq \frac{t}{2a} + \frac{r}{a}} \sin \frac{2\pi J}{t} \cos \left(\frac{2\pi jaJ}{t} - \frac{2\pi jr}{t} \right) + \mathcal{O}\left(\frac{a}{t}\right) \\
& - 2 \sum_{r \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \sum_{\frac{t}{2a} - \frac{r}{a} < J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \cos \left(\frac{2\pi jaJ}{t} + \frac{2\pi jr}{t} \right) + \mathcal{O}\left(\frac{a}{t}\right);
\end{aligned}$$

now on, we'll ignore all remainders giving a contribution already into (2): the first product here is $\ll \frac{1}{t}$. Then, from the quoted [D] above, the second ("non-sporadic") term is ($\forall j \neq \pm \bar{a}(t)$, here)

$$\mathcal{O}\left(\frac{a}{t}\right) \sum_{J \leq \frac{t}{2a}} \cos \frac{2\pi J}{t} \cos \frac{2\pi jaJ}{t} = \mathcal{O}\left(\frac{a}{t}\right) \sum_{J \leq \frac{t}{2a}} \cos \frac{2\pi jaJ}{t} + \mathcal{O}\left(\frac{1}{a^2}\right) \ll \mathbf{1}_{j=0} + \frac{a}{t} \left(\mathbf{1}_{j \neq 0} \frac{1}{\left\| \frac{ja}{t} \right\|} \right) + \frac{1}{a^2}$$

Since we have $\sin \frac{2\pi J}{t} = \frac{\pi}{a} + \mathcal{O}\left(\frac{1}{t}\right)$ into sporadic terms, these are:

$$\begin{aligned}
& -\frac{2\pi}{a} \sum_{r \leq a/2} \sin \frac{2\pi r \bar{t}}{a} \left(\sum_{\frac{t}{2a} < J \leq \frac{t}{2a} + \frac{r}{a}} \cos \left(\frac{2\pi jaJ}{t} - \frac{2\pi jr}{t} \right) + \sum_{\frac{t}{2a} - \frac{r}{a} < J \leq \frac{t}{2a}} \cos \left(\frac{2\pi jaJ}{t} + \frac{2\pi jr}{t} \right) \right) \\
& (\Rightarrow \sum_r \sin \frac{2\pi r \bar{t}}{a} \Sigma_2(0) \ll 1), \text{ i.e., applying RECIPROCITY } \frac{\bar{t}}{a} \equiv -\frac{\bar{a}}{t} + \frac{1}{at} \pmod{1} \text{ and expanding cosines,} \\
& -\frac{2\pi}{a} \cos \frac{2\pi ja}{t} \left[\frac{t}{2a} + \frac{1}{2} \right] \sum_{r \leq a/2} \left(\left[\frac{t}{2a} + \frac{r}{a} \right] - \left[\frac{t}{2a} \right] \right) \sin \frac{2\pi r \bar{a}}{t} \cos \frac{2\pi jr}{t} \\
& -\frac{2\pi}{a} \sin \frac{2\pi ja}{t} \left[\frac{t}{2a} + \frac{1}{2} \right] \sum_{r \leq a/2} \left(\left[\frac{t}{2a} + \frac{r}{a} \right] - \left[\frac{t}{2a} \right] \right) \sin \frac{2\pi r \bar{a}}{t} \sin \frac{2\pi jr}{t} \\
& -\frac{2\pi}{a} \cos \frac{2\pi ja}{t} \left[\frac{t}{2a} \right] \sum_{r \leq a/2} \left(\left[\frac{t}{2a} \right] - \left[\frac{t}{2a} - \frac{r}{a} \right] \right) \sin \frac{2\pi r \bar{a}}{t} \cos \frac{2\pi jr}{t} \\
& +\frac{2\pi}{a} \sin \frac{2\pi ja}{t} \left[\frac{t}{2a} \right] \sum_{r \leq a/2} \left(\left[\frac{t}{2a} \right] - \left[\frac{t}{2a} - \frac{r}{a} \right] \right) \sin \frac{2\pi r \bar{a}}{t} \sin \frac{2\pi jr}{t} + \mathcal{O}\left(\frac{1}{t}\right) \\
& \ll \frac{1}{a} \left(\left| \sum_{R_1 < r \leq R_2} e_t((\bar{a}+j)r) \right| + \left| \sum_{R_1 < r \leq R_2} e_t((\bar{a}-j)r) \right| + \left| \sum_{R_3 < r \leq R_4} e_t((\bar{a}+j)r) \right| + \left| \sum_{R_3 < r \leq R_4} e_t((\bar{a}-j)r) \right| \right) + \frac{1}{t} \\
& \ll \frac{a}{t} + \frac{1}{a} \left(\frac{1}{\left\| \frac{j+\bar{a}}{t} \right\|} + \frac{1}{\left\| \frac{j-\bar{a}}{t} \right\|} \right). \quad \square
\end{aligned}$$

3. Proof of the Theorem.

A kind of “elementary dispersion” ([C, Lemma 1]) gives (writing $n \sim N$ in sums for $N < n \leq 2N$), defining

$$W(a) \stackrel{\text{def}}{=} \mathbf{1}_{[-2h, 2h]}(a) \max(2h - 3|a|, |a| - 2h) \quad \text{AND} \quad \mathcal{C}_f(a) \stackrel{\text{def}}{=} \sum_{n \sim N} f(n) f(n - a),$$

$$I_f(N, h) = \sum_{a \neq 0} W(a) \mathcal{C}_f(a) + \mathcal{O}(N^\varepsilon (Nh + h^3)).$$

(We’ll refer to the first kind of remainders as the “DIAGONAL-TYPE”, while the others will be the “TAILS”). Then, after using $f(n - a) = \sum_{q|n-a} g(q)$, we organize the q -sum w.r.t. $(a, q) = \text{GCD}(a, q)$, say:

$$\sum_{a \neq 0} W(a) \mathcal{C}_f(a) = \sum_{a \neq 0} W(a) \sum_{\ell|a} \sum_{(q,a)=\ell} g(q) \sum_{\substack{n \sim N \\ n \equiv a(q)}} f(n)$$

which, changing variables and using $|a| \leq 2h$ (otherwise $W = 0$), is

$$\sum_{a \neq 0} W(a) \mathcal{C}_f(a) = \sum_{\ell \leq 2h} \sum_{a \neq 0} W(a\ell) \sum_{(q,a)=1} g(\ell q) \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv a(q)}} f(\ell m),$$

then (using $Q \ll N$ from g definition and $W(a) \ll h$ UNIFORMLY on a) we “cut” the divisors ℓ at $G = o(h)$:

$$\sum_{a \neq 0} W(a) \mathcal{C}_f(a) = \sum_{\ell \leq G} \sum_{a \neq 0} W(a\ell) \sum_{(q,a)=1} g(\ell q) \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv a(q)}} f(\ell m) + \mathcal{O}\left(N^\varepsilon \frac{Nh^2}{G}\right);$$

also, we may assume $|a| > h/G$, otherwise the error is again the same (using $h = o(N)$, here):

$$\sum_{a \neq 0} W(a) \mathcal{C}_f(a) = \sum_{\ell \leq G} \sum_{|a| > \frac{h}{G}} W(a\ell) \sum_{(q,a)=1} g(\ell q) \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv a(q)}}^* f(\ell m) + \mathcal{O}\left(N^\varepsilon \left(\frac{Nh^2}{G}\right)\right).$$

Hereon we’ll abbreviate the condition “ $(m, q) = 1$ ” with a “*” in the m -sum.

(We use coprimality of a and q to insert a limitation which is immaterial: $m \equiv a(q) \Rightarrow (m, q) = 1$.)

Since remainders are negligible, we write $\sim$ to leave them. (No confusion arises with $\sim$ in summations !)

Thus

$$I_f(N, h) \sim \sum_{\ell \leq G} \sum_{|a| > \frac{h}{G}} W(a\ell) \sum_{(q,a)=1} g(\ell q) \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv a(q)}}^* f(\ell m)$$

and, applying the orthogonality [V] of additive characters $e_q(ja) \stackrel{\text{def}}{=} e(\frac{ja}{q})$, where $e(\alpha) \stackrel{\text{def}}{=} e^{2\pi i \alpha}$, this is

$$I_f(N, h) \sim \sum_{\ell \leq G} \sum_{|a| > \frac{h}{G}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{0 < |j| \leq q/2} e_q(ja) \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) e_q(-jm)$$

since $j = 0$ terms give diagonal-type remainders, due to: $\sum_{a \neq 0} W(a\ell) \ll h$ [C, Lemma 4, (1)]. Being W EVEN

$$\begin{aligned} I_f(N, h) &\sim \sum_{\ell \leq G} \sum_{|a| > \frac{h}{G}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{0 < |j| \leq \frac{q}{2}} \cos \frac{2\pi ja}{q} \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) \cos \frac{2\pi jm}{q} \\ &\sim 2 \sum_{\ell \leq G} \sum_{\frac{h}{G} < a \leq \frac{2h}{\ell}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{0 < |j| \leq \frac{q}{2}} \cos \frac{2\pi ja}{q} \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) \cos \frac{2\pi jm}{q} \end{aligned}$$

and, now, we exploit the coprimality of a with q , changing variable:

$$I_f(N, h) \sim 2 \sum_{\ell \leq G} \sum_{\frac{h}{G} < a \leq \frac{2h}{\ell}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{0 < |j| \leq q/2} \cos \frac{2\pi j}{q} \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) \cos \frac{2\pi j m \bar{a}}{q}$$

We rearrange, then, the terms, depending on residue classes modulo q :

$$I_f(N, h) \sim 2 \sum_{\ell \leq G} \sum_{\frac{h}{G} < a \leq \frac{2h}{\ell}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{\substack{0 < |j|, |k| \leq q/2 \\ (j,q)=(k,q)}} \cos \frac{2\pi j}{q} \cos \frac{2\pi k \bar{a}}{q} \sum_{\substack{m \sim \frac{N}{\ell} \\ jm \equiv k(q)}}^* f(\ell m)$$

Therefore, applying RECIPROCITY $\frac{\bar{a}}{q} \equiv -\frac{\bar{q}}{a} + \frac{1}{aq} \pmod{1}$, we separate into two sums, say $I_f \sim S_1 + S_2$:

$$S_1 \stackrel{def}{=} 2 \sum_{\ell \leq G} \sum_{\frac{h}{G} < a \leq \frac{2h}{\ell}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \sum_{0 < |j|, |k| \leq q/2} \cos \frac{2\pi j}{q} \cos \frac{2\pi k \bar{q}}{a} \sum_{\substack{m \sim \frac{N}{\ell} \\ jm \equiv k(q)}}^* f(\ell m)$$

and

$$S_2 \stackrel{def}{=} 2 \sum_{\ell \leq G} \sum_{\frac{h}{G} < a \leq \frac{2h}{\ell}} W(a\ell) \sum_{(q,a)=1} \frac{g(\ell q)}{q} \Sigma,$$

where, say,

$$\Sigma \stackrel{def}{=} \sum_{\substack{0 < |j|, |k| \leq q/2 \\ (j,q)=(k,q)}} \cos \frac{2\pi j}{q} \left(\cos \frac{2\pi k \bar{q}}{a} \left(\cos \frac{2\pi k}{aq} - 1 \right) + \sin \frac{2\pi k \bar{q}}{a} \sin \frac{2\pi k}{aq} \right) \sum_{\substack{m \sim \frac{N}{\ell} \\ jm \equiv k(q)}}^* f(\ell m)$$

We'll treat first S_2 , as it contains “remainder terms”, here. Collect all j with same (j, q) and change j, k :

$$\begin{aligned} \Sigma &= \sum_{\substack{t|q \\ t < q}} \sum_{\substack{0 < |j|, |k| \leq \frac{q}{2t} \\ (j, \frac{q}{t})=(k, \frac{q}{t})=1}} \cos \frac{2\pi j}{q/t} \left(\cos \frac{2\pi k \bar{q}/t}{a} \left(\cos \frac{2\pi k}{a(q/t)} - 1 \right) + \sin \frac{2\pi k \bar{q}/t}{a} \sin \frac{2\pi k}{a(q/t)} \right) \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv \bar{j}k(q/t)}}^* f(\ell m) \stackrel{F}{=} \\ &\stackrel{F}{=} \sum_{\substack{t|q \\ t > 1}} \sum_{\substack{0 < |k| \leq \frac{q}{2t} \\ (k, t)=1}} \left(\cos \frac{2\pi k \bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k \bar{t}}{a} \sin \frac{2\pi k}{at} \right) \sum_{\substack{0 < |j| \leq \frac{q}{2t} \\ (j, t)=1}} \cos \frac{2\pi j}{t} \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv \bar{j}k(t)}}^* f(\ell m) \end{aligned}$$

where “F” means we “flipped” the divisors t . Our inner sum over j, m is, again through orthogonality,

$$\begin{aligned} \sum_{\substack{0 < |j| \leq \frac{q}{2t} \\ (j, t)=1}} \cos \frac{2\pi j}{t} \sum_{\substack{m \sim \frac{N}{\ell} \\ m \equiv \bar{j}k(t)}}^* f(\ell m) &= \frac{1}{t} \sum_{\substack{0 < |j| \leq \frac{q}{2t} \\ (j, t)=1}} \cos \frac{2\pi j}{t} \sum_{0 \leq |s| \leq t/2} e_t(sk \bar{j}) \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) e_t(-sm) \\ &= \frac{\mu(t)}{t} \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) + \frac{1}{2t} \sum_{0 < |s| \leq t/2} [S(1, sk; t) + S(-1, sk; t)] \sum_{m \sim \frac{N}{\ell}}^* f(\ell m) e_t(-sm), \end{aligned}$$

where μ is Möbius function (see above, in the introduction),

$$c_t(n) \stackrel{def}{=} \sum_{j \leq t, (j, t)=1} e_t(jn) = \varphi(t) \frac{\mu(t/(t, n))}{\varphi(t/(t, n))} = \sum_{\substack{0 < |j| \leq \frac{q}{2t} \\ (j, t)=1}} \cos \frac{2\pi j n}{t} \quad \text{is RAMANUJAN SUM}$$

($n = 1$ gives $\mu(t)$ here), see quoted [D], with Euler function $\varphi(n) \stackrel{def}{=} \sum_{j \leq n, (j,n)=1} 1$ and

$$S(a, b; c) \stackrel{def}{=} \sum_{\substack{j \leq c, (j,c)=1 \\ j\bar{j} \equiv 1(c)}} e_c(ja + \bar{j}b) \quad \text{is KLOOSTERMAN SUM}$$

that can be estimated using Weil-Esternmann bound, see [I-K, chap.11],

$$S(a, b; c) \ll d(t) \sqrt{(a, b, c)} \sqrt{c} \quad (\Rightarrow [S(1, ks; t) + S(-1, ks; t)] \ll \sqrt{t}, \text{ here}).$$

We'll use this bound in a while. First, see that (recall $s \not\equiv 0(t)$ and use $\mathbf{1}_{(m,q)=1} = \sum_{d|q, d|m} \mu(d)$, [T])

$$\begin{aligned} \sum_{m \sim \frac{N}{t}}^* f(\ell m) e_t(-sm) &= \sum_{q'} g(q') \sum_{\substack{n \sim N \\ n \equiv 0(q'), n \equiv 0(\ell) \\ (\frac{N}{t}, q)=1}} e_t(-s(n/\ell)) \\ &= \sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'} q)=1}} g(\ell' q'') \sum_{\substack{m \sim N/(\ell q'') \\ (m, q)=1}} e_t(-sq'' m) = \sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'} q)=1}} g(\ell' q'') \sum_{d|q} \mu(d) \sum_{m \sim N/(\ell q'' d)} e_t(-sq'' dm) \\ &= \sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'} q)=1}} g(\ell' q'') \sum_{\substack{d|q \\ sq'' d \equiv 0(t)}} \mu(d) \left(\frac{N}{\ell q'' d} + \mathcal{O}(1) \right) + \mathcal{O} \left(N^\varepsilon \sum_{\ell'|\ell} \sum_{\substack{q'' \sim \frac{Q}{\ell'} \\ (q'', \frac{\ell}{\ell'} q)=1}} \sum_{\substack{d|q \\ sq'' d \not\equiv 0(t)}} \frac{1}{\left\| \frac{sq'' d}{t} \right\|} \right) \end{aligned}$$

(we used, when $\|\beta\| \neq 0$, the classical bound $\sum_{N_1 \leq n \leq N_2} e(n\beta) \ll 1/\|\beta\|$, see quoted [D]) gives

$$\begin{aligned} (*) \quad \sum_{(q,a)=1} \frac{g(\ell q)}{q} \Sigma &= \sum_{\substack{1 < t \leq \frac{2Q}{t} \\ (t,a)=1}} \left(\sum_{(q,a)=1} \frac{g(\ell t q)}{q} \right) \frac{1}{t} \sum_{\substack{0 < |k| \leq \frac{t}{2} \\ (k,t)=1}} \left(\cos \frac{2\pi k \bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k \bar{t}}{a} \sin \frac{2\pi k}{at} \right) \times \\ &\times \left(\frac{\mu(t)}{t} \sum_{\substack{m \sim \frac{N}{t} \\ (m,qt)=1}} f(\ell m) + \frac{1}{2t} \sum_{0 < |s| \leq t/2} [S(1, sk; t) + S(-1, sk; t)] \left[\sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'} qt)=1}} g(\ell' q'') \sum_{\substack{d|qt \\ sq'' d \equiv 0(t)}} \mu(d) \times \right. \right. \\ &\left. \left. \times \left(\frac{N}{\ell q'' d} + \mathcal{O}(1) \right) + \mathcal{O} \left(N^\varepsilon \sum_{\ell'|\ell} \sum_{\substack{q'' \sim \frac{Q}{\ell'} \\ (q'', \frac{\ell}{\ell'} qt)=1}} \sum_{\substack{d|qt \\ sq'' d \not\equiv 0(t)}} \frac{1}{\left\| \frac{sq'' d}{t} \right\|} \right) \right] \right) \end{aligned}$$

which we'll distinguish into $t \leq \frac{T}{\ell}$ and $t > \frac{T}{\ell}$; for the former

$$\begin{aligned} \sum_{\substack{1 < t \leq \frac{T}{\ell} \\ (t,a)=1}} \left(\sum_{(q,a)=1} \frac{g(\ell t q)}{q} \right) \frac{1}{t} \sum_{\substack{0 < |k| \leq \frac{t}{2} \\ (k,t)=1}} \left(\cos \frac{2\pi k \bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k \bar{t}}{a} \sin \frac{2\pi k}{at} \right) \times \\ \times \left(\frac{\mu(t)}{t} \sum_{\substack{m \sim \frac{N}{t} \\ (m,qt)=1}} f(\ell m) + \frac{1}{2t} \sum_{0 < |s| \leq t/2} [S(1, sk; t) + S(-1, sk; t)] \left[\sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'} qt)=1}} g(\ell' q'') \sum_{\substack{d|qt \\ sq'' d \equiv 0(t)}} \mu(d) \mathcal{O}(1) \right. \right. \end{aligned}$$

$$+ \mathcal{O} \left(N^\varepsilon \sum_{\ell'|\ell} \sum_{\substack{q'' \sim \frac{Q}{\ell'} \\ (q'', \frac{\ell}{\ell'}qt)=1}} \sum_{\substack{d|qt \\ sq''d \not\equiv 0(t)}} \frac{1}{\left\| \frac{sq''d}{t} \right\|} \right) \ll \frac{N}{a\ell} + \frac{QT^{3/2}}{a\ell^{3/2}} + \frac{T^{5/2}}{a\ell^{5/2}},$$

having used the trivial bound $\frac{1}{t} \sum_{\substack{0 < |k| \leq \frac{t}{2} \\ (k, t)=1}} \left(\cos \frac{2\pi k\bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k\bar{t}}{a} \sin \frac{2\pi k}{at} \right) \ll \frac{1}{a}$ and, esp.,

$$\begin{aligned} & \sum_{\substack{1 < t \leq \frac{T}{\ell} \\ (t, a)=1}} \left(\sum_{(q, a)=1} \frac{g(\ell tq)}{q} \right) \frac{1}{t} \sum_{\substack{0 < |k| \leq \frac{t}{2} \\ (k, t)=1}} \left(\cos \frac{2\pi k\bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k\bar{t}}{a} \sin \frac{2\pi k}{at} \right) \times \\ & \times \frac{1}{2t} \sum_{0 < |s| \leq t/2} [S(1, sk; t) + S(-1, sk; t)] \mathcal{O} \left(N^\varepsilon \sum_{\ell'|\ell} \sum_{\substack{q'' \sim \frac{Q}{\ell'} \\ (q'', \frac{\ell}{\ell'}qt)=1}} \sum_{\substack{d|qt \\ sq''d \not\equiv 0(t)}} \frac{1}{\left\| \frac{sq''d}{t} \right\|} \right) \\ & \ll \sum_{t \leq \frac{T}{\ell}} \frac{1}{a} \frac{1}{t} \sum_{s \leq \frac{t}{2}} \sqrt{t} \sum_{\ell'|\ell} \sum_{q'' \sim \frac{Q}{\ell'}} \sum_{d \leq \frac{2Q}{\ell'}} \frac{1}{d} \sum_{\substack{r \leq t/2 \\ sq''d \equiv \pm r(t)}} \frac{t}{r} \ll \sum_{t \leq \frac{T}{\ell}} \frac{\sqrt{t}}{a} \sum_{t'|t} \left(\frac{Q}{t/t'} + 1 \right) t \sum_{r \leq \frac{t}{2t'}} \frac{1}{t'r} \ll \frac{QT^{\frac{3}{2}}}{a\ell^{\frac{3}{2}}} + \frac{T^{\frac{5}{2}}}{a\ell^{\frac{5}{2}}}. \end{aligned}$$

Here we use the Weil bound for Kloosterman sums. Last term of $t \leq T/\ell$ is

$$\begin{aligned} & \sum_{\substack{1 < t \leq \frac{T}{\ell} \\ (t, a)=1}} \left(\sum_{(q, a)=1} \frac{g(\ell tq)}{q} \right) \frac{1}{t} \sum_{\substack{0 < |k| \leq \frac{t}{2} \\ (k, t)=1}} \left(\cos \frac{2\pi k\bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k\bar{t}}{a} \sin \frac{2\pi k}{at} \right) \times \\ & \times \frac{1}{2t} \sum_{0 < |s| \leq t/2} [S(1, sk; t) + S(-1, sk; t)] \frac{N}{\ell} \sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'}qt)=1}} \frac{g(\ell'q'')}{q''} \sum_{\substack{d|qt \\ sq''d \equiv 0(t)}} \frac{\mu(d)}{d} \ll \frac{N}{a\ell} \end{aligned}$$

since terms after $\times$ can be treated opening the Kloosterman sum, see the above, becoming

$$\frac{N}{\ell t} \sum_{(j, t)=1} \cos \frac{2\pi j}{t} \sum_{t'|t} \sum_{\ell'|\ell} \sum_{\substack{q'' \\ (q'', \frac{\ell}{\ell'}qt)=1}} \frac{g(\ell'q'')}{q''} \sum_{\substack{d|qt \\ (d, t)=t'}} \frac{\mu(d)}{d} \sum_{\substack{0 < |s| \leq t/2 \\ s \equiv 0(t/t')}} e_t(k\bar{j}s),$$

but this time we gain t , from $\sum_{(j, t)=1} \cos \frac{2\pi j}{t} = \mu(t)$ (due to independence of the following sum from j):

$$\sum_{\substack{0 < |s| \leq t/2 \\ s \equiv 0(t/t')}} e_t(k\bar{j}s) = \sum_{0 < |s| \leq t'/2} e_{t'}(k\bar{j}s) = \mathbf{1}_{k \equiv 0(t')} - \frac{1}{t'}.$$

Finally, the $t \leq T/\ell$ terms of $(*)$ are in total $\ll \frac{N}{a\ell} + \frac{QT^{3/2}}{a\ell^{3/2}} + \frac{T^{5/2}}{a\ell^{5/2}}$; contribute to S_2 as $\ll (Q + T)T^{3/2}h$.

We pass to the terms $t > T/\ell$.

For these (hence, t is “LARGE”), we use Lemma 1:

$$\sum_{\substack{t|q \\ t \text{ LARGE}}} \sum_{\substack{0 < |j| \leq \frac{t}{2} \\ (j, t)=1}} \sum_{0 < |k| \leq \frac{t}{2}} \left[\cos \frac{2\pi k\bar{t}}{a} \left(\cos \frac{2\pi k}{at} - 1 \right) + \sin \frac{2\pi k\bar{t}}{a} \sin \frac{2\pi k}{at} \right] \cos \frac{2\pi jk}{t} \sum_{\substack{m \sim \frac{N}{t} \\ m \equiv \bar{j}(t)}}^* f(\ell m)$$

$$= -2 \sum_{\substack{t|q \\ t \text{ LARGE}}} \sum_{\substack{|j| \leq \frac{t}{2} \\ (j,t)=1}} \sum_{|r| \leq \frac{t}{2}} \sin \frac{2\pi r \bar{t}}{a} \sin \frac{2\pi jr}{t} \sum_{J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \sin \frac{2\pi jaJ}{t} \sum_{\substack{m \sim \frac{N}{t} \\ jm \equiv 1(t)}}^* f(\ell m) + \mathcal{O} \left(N^\varepsilon \sum_{\substack{t|q \\ t \text{ LARGE}}} \left(\frac{1}{a} + \frac{a}{t} \right) \frac{N}{\ell} \right)$$

so to apply quoted [D] $\sum_{N_1 < n \leq N_2} e(\alpha n) \ll \min(N_2 - N_1, \frac{1}{\|\alpha\|})$ and partial summation (with RECIPROCITY)

$$\begin{aligned} & \sum_{\substack{0 < |j| \leq \frac{t}{2} \\ j \neq \pm \bar{a}(t)}} \left| \sum_{|r| \leq \frac{t}{2}} \sin \frac{2\pi r \bar{t}}{a} \sin \frac{2\pi jr}{t} \right| \left| \sum_{J \leq \frac{t}{2a}} \sin \frac{2\pi J}{t} \sin \frac{2\pi jaJ}{t} \right| \ll \frac{1}{a} \sum_{\substack{0 < |j| \leq \frac{t}{2} \\ j \neq \pm \bar{a}(t)}} \min \left(a, \frac{1}{\left\| \frac{j \pm \bar{a}}{t} \right\|} \right) \min \left(\frac{t}{a}, \frac{1}{\left\| \frac{ja}{t} \right\|} \right) \\ & \ll \frac{1}{a} \sum_{1 < r \leq \frac{t}{2}} \left(\sum_{j \equiv \pm \bar{a}r(t)} 1 \right) \min \left(a, \frac{1}{\left\| \frac{(r \pm 1)\bar{a}}{t} \right\|} \right) \min \left(\frac{t}{a}, \frac{t}{r} \right) \ll \frac{1}{a} \sum_{r < a} \frac{t}{a} \min \left(a, \frac{1}{\left\| \frac{r\bar{a}}{t} \right\|} \right) + \frac{t}{a} + \frac{1}{a} \sum_{j \leq \frac{t}{2a}} \frac{t}{j} \\ & + \frac{1}{a} \sum_{\substack{a < r \leq \frac{t}{2} \\ r \neq 0(a)}} \frac{t}{r} \min \left(a, \frac{1}{\left\| \frac{r\bar{a}}{t} \right\|} \right) \ll \frac{1}{a} \sum_{0 < |r| \leq \frac{t}{2}} \frac{t}{a} \min \left(a, \frac{1}{\left\| \frac{r\bar{t}}{a} \right\| - \frac{1}{2a}} \right) + \frac{1}{a} \sum_{\substack{a < r \leq \frac{t}{2} \\ r \neq 0(a)}} \frac{t}{r} \min \left(a, \frac{1}{\left\| \frac{r\bar{t}}{a} \right\| - \frac{|r|}{at}} \right) + \frac{t}{a} \log t \\ & \ll \frac{1}{a} \left(\frac{t}{a} \sum_{0 < |r| \leq \frac{t}{2}} \min \left(a, \frac{1}{\left| \frac{r}{a} - \frac{1}{2a} \right|} \right) + t \sum_{j \leq \frac{t}{2a}} \frac{1}{ja} \sum_{0 < |r-j| \leq \frac{t}{2}} \min \left(a, \frac{1}{\left| \frac{r-j}{a} - \frac{1}{2a} \right|} \right) + t \log t \right) \ll \frac{t}{a}. \end{aligned}$$

Here we may assume $j \neq \pm \bar{a}(t)$, since the corresponding terms give a contribute already in those before:

$$\lll \sum_{t|q, t \text{ LARGE}} \left| \sum_{|r| \leq \frac{t}{2}} \sin \frac{2\pi r \bar{t}}{a} \sin \frac{2\pi \bar{a}r}{t} \right| \left| \sum_{J \leq \frac{t}{2a}} \sin^2 \frac{2\pi J}{t} \right| \frac{N}{\ell t} \lll \sum_{t|q, t \text{ LARGE}} \frac{N}{a\ell}.$$

All these terms and the previous (with t LARGE, say) give (using $\sum_q \frac{g(\ell tq)}{q} \lll 1$)

$$\lll \sum_{T/\ell < t \leq 2Q/\ell} \frac{1}{t} \left(\frac{1}{a} + \frac{a}{t} \right) \frac{N}{\ell} \lll \frac{N}{\ell a} + \frac{Na}{T}$$

contributes to the LARGE t in (*). The first is diagonal-type, the others make $\lll Nh^3/T$ for S_2 .

Thus, in all, the sum $S_2 \lll Nh + (Q + T)T^{3/2}h + Nh^3/T$ and (here T is somehow small respect to Q) we choose $T = (Nh^2/Q)^{2/5}$ (optimally). Also, this gives us $h = o(T)$, from $Q = o(N/\sqrt{h})$.

As regards S_1 , we may write, with $A_1(x) \stackrel{def}{=} x, \forall x \in [-1/2, 1/2]$ (ODD and) 1-PERIODICIZED

$$\begin{aligned} & \sum_j' \cos \frac{2\pi j}{q} \sum_k' \cos \frac{2\pi k \bar{q}}{a} \sum_{jm \equiv k(q)}^* f(\ell m) = \sum_j' \cos \frac{2\pi j}{q} \sum_{m \sim \frac{N}{t}}^* f(\ell m) \cos \left(\frac{2\pi \bar{q}}{a} q A_1 \left(\frac{jm}{q} \right) \right) \\ & = \sum_j' \cos \frac{2\pi j}{q} \sum_{m \sim \frac{N}{t}}^* f(\ell m) \cos \left(\frac{2\pi}{a} A_1 \left(\frac{jm}{q} \right) \right) = \sum_j' \cos \frac{2\pi j}{aq} \sum_{m \sim \frac{N}{t}}^* f(\ell m) \cos \frac{2\pi j \bar{m}}{q} \end{aligned}$$

where $'$ means: in $[-q/2, q/2]$ and not zero. Here $0 < |k| \leq q/2$ and $k \equiv jm(q)$ force k to be $qA_1(jm/q)$. Then,

$$\sum_{0 < |j| \leq \frac{q}{2}} \cos \frac{2\pi j}{aq} \sum_{m \sim \frac{N}{t}}^* f(\ell m) \cos \frac{2\pi j \bar{m}}{q} \ll \frac{1}{a} \sum_{m \sim \frac{N}{t}}^* |f(\ell m)| \frac{1}{\left\| \frac{\bar{m}}{q} \right\|} \lll \frac{1}{a} \sum_{m \sim \frac{N}{t}}^* \frac{1}{\left\| \frac{\bar{m}}{q} \right\|}$$

using quoted [D] and this $\Rightarrow S_1$ of DIAGONAL-TYPE:

$$S_1 \lll Nh. \quad \square$$

References

- [C] Coppola, G.- *On the correlations, Selberg integral and symmetry of sieve functions in almost all short intervals* - <http://arxiv.org/abs/0709.3648v3> - 9pp. (electronic).
- [D] Davenport, H.- *Multiplicative Number Theory* - Third Edition, GTM 74, Springer, New York, 2000. MR 2001f:11001
- [I-K] Iwaniec, H. and Kowalski, E.- *Analytic Number Theory* - American Mathematical Society Colloquium Publications, 53. AMS, Providence, RI, 2004. xii+615 pp. ISBN: 0-8218-3633-1 MR 2005h:11005
- [T] Tenenbaum, G. - *Introduction to analytic and probabilistic number theory* - Cambridge studies in advanced mathematics : 46, Cambridge University Press, 1995. MR 97e:11005b
- [V] Vinogradov, I.M.- *The Method of Trigonometrical Sums in the Theory of Numbers* - Interscience Publishers LTD, London, 1954. MR 15,941b

Dr.Giovanni Coppola
DIIMA - Università degli Studi di Salerno
84084 Fisciano (SA) - ITALY
e-mail : gcoppola@diima.unisa.it